

**LOGISTIC REGRESSION**

David Kauchak  
CS158 – Fall 2019

### Admin

Assignment 6: what did you find?

Assignment 7

### Priors

Coin1 data: 3 Heads and 1 Tail  
Coin2 data: 30 Heads and 10 tails  
Coin3 data: 2 Tails  
Coin4 data: 497 Heads and 503 tails

If someone asked you what the probability of heads was for each of these coins, what would you say?

### Basic steps for probabilistic modeling

Step 1: pick a model	<b>Probabilistic models</b> Which model do we use, i.e. how do we calculate $p(\text{feature}, \text{label})$ ?
Step 2: figure out how to estimate the probabilities for the model	How do train the model, i.e. how to we we <b>estimate the probabilities</b> for the model?
Step 3 (optional): deal with overfitting	How do we deal with overfitting?

## Training revisited

What we're really doing during training is selecting the  $\theta$  that maximizes:

$$p(\theta | data)$$

i.e.

$$\theta = \operatorname{argmax}_{\theta} p(\theta | data)$$

That is we pick the most likely model parameters given the data

## Estimating revisited

We want to incorporate a prior belief of what the probabilities might be

To do this, we need to break down our probability

$$p(\theta | data) = ?$$

(Hint: Bayes rule)

## Estimating revisited

What are each of these probabilities?

$$p(\theta | data) = \frac{p(data | \theta)p(\theta)}{p(data)}$$

## Priors

likelihood of the data  
under the model

probability of different parameters,  
call the **prior**

$$p(\theta | data) = \frac{p(data | \theta)p(\theta)}{p(data)}$$

probability of seeing the data  
(regardless of model)

### Priors

$$\theta = \operatorname{argmax}_{\theta} \frac{p(\text{data} | \theta)p(\theta)}{p(\text{data})}$$

Does  $p(\text{data})$  matter for the argmax?

### Priors

likelihood of the data under the model

probability of different parameters, call the **prior**

$$\theta = \operatorname{argmax}_{\theta} p(\text{data} | \theta)p(\theta)$$

What does MLE assume for a prior on the model parameters?

### Priors

likelihood of the data under the model

probability of different parameters, call the **prior**

$$\theta = \operatorname{argmax}_{\theta} p(\text{data} | \theta)p(\theta)$$

- Assumes a **uniform prior**, i.e. all  $\Theta$  are equally likely!
- Relies solely on the likelihood

### A better approach

$$\theta = \operatorname{argmax}_{\theta} p(\text{data} | \theta)p(\theta)$$

$$\text{likelihood}(\text{data}) = \prod_{i=1}^n p_{\theta}(x_i)$$

We can use any distribution we'd like.

This allows us to impart addition **bias** into the model

### Another view on the prior

Remember, the max is the same if we take the log:

$$\theta = \operatorname{argmax}_{\theta} \log(p(\text{data} | \theta)) + \log(p(\theta))$$

$$\text{log-likelihood} = \sum_{i=1}^n \log(p(x_i))$$

We can use any distribution we'd like.  
This allows us to impart addition **bias** into the model

Does this look like something we've seen before?

### Regularization vs prior

$$\theta = \operatorname{argmax}_{\theta} \log(p(\text{data} | \theta)) + \log(p(\theta))$$


fit { likelihood based on the data / loss function based on the data } prior regularizer } model bias

$$\operatorname{argmin}_{w,b} \sum_{i=1}^n \text{loss}(yy') + \lambda \text{regularizer}(w)$$

### Prior for NB

$$\theta = \operatorname{argmax}_{\theta} \log(p(\text{data} | \theta)) + \log(p(\theta))$$

Uniform prior



$$p(x_i | y) = \frac{\text{count}(x_i, y)}{\text{count}(y)}$$

Dirichlet prior



lambda = 0 increasing

$$p(x_i | y) = \frac{\text{count}(x_i, y) + \lambda}{\text{count}(y) + \text{possible\_values\_of\_}x_i * \lambda}$$

### Prior: another view

$$p(x_1, x_2, \dots, x_m, y) = p(y) \prod_{j=1}^m p(x_j | y)$$

$$\text{MLE: } p(x_i | y) = \frac{\text{count}(x_i, y)}{\text{count}(y)}$$

What happens to our likelihood if, for one of the labels, we never saw a particular feature?

Goes to 0!

### Prior: another view

$$p(x_i | y) = \frac{\text{count}(x_i, y)}{\text{count}(y)}$$

↓

$$p(x_i | y) = \frac{\text{count}(x_i, y) + \lambda}{\text{count}(y) + \text{possible\_values\_of\_}x_i * \lambda}$$

Adding a prior avoids this!

### Smoothing

training data

$$p(x_i | y) = \frac{\text{count}(x_i, y)}{\text{count}(y)}$$

↓

$$p(x_i | y) = \frac{\text{count}(x_i, y) + \lambda}{\text{count}(y) + \text{possible\_values\_of\_}x_i * \lambda}$$

for each label, pretend like we've seen each feature value occur in  $\lambda$  additional examples

Sometimes this is also called **smoothing** because it is seen as smoothing or interpolating between the MLE and some other distribution

### Basic steps for probabilistic modeling

<p>Step 1: pick a model</p> <hr/> <p>Step 2: figure out how to estimate the probabilities for the model</p> <hr/> <p>Step 3 (optional): deal with overfitting</p>	<p><b>Probabilistic models</b></p> <p>Which model do we use, i.e. how do we calculate <math>p(\text{feature}, \text{label})</math>?</p> <p>How do train the model, i.e. how to we we <b>estimate the probabilities</b> for the model?</p> <p>How do we deal with overfitting?</p>
---	---

### Joint models vs conditional models

We've been trying to model the joint distribution (i.e. the data generating distribution):

$$p(x_1, x_2, \dots, x_m, y)$$

However, if all we're interested in is classification, why not directly model the conditional distribution:

$$p(y | x_1, x_2, \dots, x_m)$$

## A first try: linear

$$p(y | x_1, x_2, \dots, x_m) = x_1 w_1 + w_2 x_2 + \dots + w_m x_m + b$$

Any problems with this?

- Nothing constrains it to be a probability
- Could still have combination of features and weight that exceeds 1 or is below 0

## The challenge

$$x_1 w_1 + w_2 x_2 + \dots + w_m x_m + b$$

$$p(y | x_1, x_2, \dots, x_m)$$

Linear model

probability



We like linear models!

Can we transform the probability into a function that ranges over all values?

## Odds ratio

Rather than predict the probability, we can predict the ratio of 1/0 (positive/negative)

Predict the **odds** that it is 1 (true): How much more likely is 1 than 0.

Does this help us?

$$\frac{P(1 | x_1, x_2, \dots, x_m)}{P(0 | x_1, x_2, \dots, x_m)} = \frac{P(1 | x_1, x_2, \dots, x_m)}{1 - P(1 | x_1, x_2, \dots, x_m)} = x_1 w_1 + w_2 x_2 + \dots + w_m x_m + b$$

## Odds ratio

$$x_1 w_1 + w_2 x_2 + \dots + w_m x_m + b$$

$$\frac{P(1 | x_1, x_2, \dots, x_m)}{1 - P(1 | x_1, x_2, \dots, x_m)}$$

Linear model

odds ratio






Where is the dividing line between class 1 and class 0 being selected?

### Odds ratio

$$\frac{P(1|x_1, x_2, \dots, x_m)}{1 - P(1|x_1, x_2, \dots, x_m)} > \frac{P(0|x_1, x_2, \dots, x_m)}{1 - P(0|x_1, x_2, \dots, x_m)}$$

We're trying to find some transformation that transforms the odds ratio to a number that is  $-\infty$  to  $+\infty$

Does this suggest another transformation?



### Log odds (logit function)

$$x_1 w_1 + w_2 x_2 + \dots + w_m x_m + b = \log \frac{P(1|x_1, x_2, \dots, x_m)}{1 - P(1|x_1, x_2, \dots, x_m)}$$

Linear regression

odds ratio



How do we get the probability of an example?

### Log odds (logit function)

$$\log \frac{P(1|x_1, x_2, \dots, x_m)}{1 - P(1|x_1, x_2, \dots, x_m)} = w_1 x_1 + w_2 x_2 + \dots + w_m x_m + b$$

$$\frac{P(1|x_1, x_2, \dots, x_m)}{1 - P(1|x_1, x_2, \dots, x_m)} = e^{w_1 x_1 + w_2 x_2 + \dots + w_m x_m + b}$$

$$P(1|x_1, x_2, \dots, x_m) = (1 - P(1|x_1, x_2, \dots, x_m)) e^{w_1 x_1 + w_2 x_2 + \dots + w_m x_m + b}$$


...

$$P(1|x_1, x_2, \dots, x_m) = \frac{1}{1 + e^{-(w_1 x_1 + w_2 x_2 + \dots + w_m x_m + b)}}$$

anyone recognize this?

## Logistic function

$$\text{logistic} = \frac{1}{1 + e^{-x}}$$



## Logistic regression

How would we classify examples once we had a trained model?

$$\log \frac{P(1|x_1, x_2, \dots, x_m)}{1 - P(1|x_1, x_2, \dots, x_m)} = w_1 x_1 + w_2 x_2 + \dots + w_m x_m + b$$

If the sum  $> 0$  then  $p(1)/p(0) > 1$ , so positive

if the sum  $< 0$  then  $p(1)/p(0) < 1$ , so negative

Still a *linear* classifier (decision boundary is a line)

## Training logistic regression models

How should we learn the parameters for logistic regression (i.e. the  $w$ 's and  $b$ )?

$$\log \frac{P(1|x_1, x_2, \dots, x_m)}{1 - P(1|x_1, x_2, \dots, x_m)} = w_1 x_1 + w_2 x_2 + \dots + w_m x_m + b$$

parameters

$$P(1|x_1, x_2, \dots, x_m) = \frac{1}{1 + e^{-(w_1 x_1 + w_2 x_2 + \dots + w_m x_m + b)}}$$

## MLE logistic regression

Find the parameters that maximize the likelihood (or log-likelihood) of the data:

$$\begin{aligned} \text{log-likelihood} &= \sum_{i=1}^n \log(p(x_i)) \\ &= \sum_{i=1}^n \log\left(\frac{1}{1 + e^{-y_i(w_1 x_{i1} + w_2 x_{i2} + \dots + w_m x_{im} + b)}}\right) \quad \text{assume labels } 1, -1 \\ &= \sum_{i=1}^n -\log(1 + e^{-y_i(w_1 x_{i1} + w_2 x_{i2} + \dots + w_m x_{im} + b)}) \end{aligned}$$



## MLE logistic regression

$$\text{log-likelihood} = \sum_{i=1}^n -\log(1 + e^{-y_i(w_1x_{i1} + w_2x_{i2} + \dots + w_mx_{im} + b)})$$

We want to maximize, i.e.

$$\begin{aligned} MLE(\text{data}) &= \operatorname{argmax}_{w,b} \text{log-likelihood}(\text{data}) \\ &= \operatorname{argmax}_{w,b} \sum_{i=1}^n -\log(1 + e^{-y_i(w_1x_{i1} + w_2x_{i2} + \dots + w_mx_{im} + b)}) \\ &= \operatorname{argmin}_{w,b} \sum_{i=1}^n \log(1 + e^{-y_i(w_1x_{i1} + w_2x_{i2} + \dots + w_mx_{im} + b)}) \end{aligned}$$

Look familiar? Hint: anybody read the book?

## MLE logistic regression

$$\operatorname{argmin}_{w,b} \sum_{i=1}^n \log(1 + e^{-y_i(w_1x_{i1} + w_2x_{i2} + \dots + w_mx_{im} + b)})$$

Surrogate loss functions:

Zero/one:  $\ell^{(0/1)}(y, \hat{y}) = \mathbf{1}[y \neq \hat{y}]$

Hinge:  $\ell^{(\text{hin})}(y, \hat{y}) = \max\{0, 1 - y\hat{y}\}$

Logistic:  $\ell^{(\text{log})}(y, \hat{y}) = \frac{1}{\log 2} \log(1 + \exp[-y\hat{y}])$

Exponential:  $\ell^{(\text{exp})}(y, \hat{y}) = \exp[-y\hat{y}]$

Squared:  $\ell^{(\text{sq})}(y, \hat{y}) = (y - \hat{y})^2$

## logistic regression: three views

$$\log \frac{P(1|x_1, x_2, \dots, x_m)}{1 - P(1|x_1, x_2, \dots, x_m)} = w_0 + w_1x_1 + w_2x_2 + \dots + w_mx_m \quad \text{linear classifier}$$

$$P(1|x_1, x_2, \dots, x_m) = \frac{1}{1 + e^{-(w_0 + w_1x_1 + w_2x_2 + \dots + w_mx_m)}} \quad \text{conditional model logistic}$$

$$\operatorname{argmin}_{w,b} \sum_{i=1}^n \log(1 + e^{-y_i(w_1x_{i1} + w_2x_{i2} + \dots + w_mx_{im} + b)}) \quad \text{linear model minimizing logistic loss}$$

## Overfitting

$$\operatorname{argmin}_{w,b} \sum_{i=1}^n \log(1 + e^{-y_i(w_1x_{i1} + w_2x_{i2} + \dots + w_mx_{im} + b)})$$

If we minimize this loss function, in practice, the results aren't great and we tend to overfit

Solution?

### Regularization/prior

---

$$\operatorname{argmin}_{w,b} \sum_{i=1}^n \log(1 + e^{-y_i(w_1x_{i1} + w_2x_{i2} + \dots + w_mx_{im} + b)}) + \lambda \operatorname{regularizer}(w,b)$$

or

$$\operatorname{argmin}_{w,b} \sum_{i=1}^n \log(1 + e^{-y_i(w_1x_{i1} + w_2x_{i2} + \dots + w_mx_{im} + b)}) - \log(p(w,b))$$

What are some of the regularizers we know?

### Regularization/prior


---

L2 regularization:

$$\operatorname{argmin}_{w,b} \sum_{i=1}^n \log(1 + e^{-y_i(w_1x_{i1} + w_2x_{i2} + \dots + w_mx_{im} + b)}) + \lambda \|w\|^2$$

Gaussian prior:

$p(w,b) \sim$



### Regularization/prior

---

L2 regularization:

$$\operatorname{argmin}_{w,b} \sum_{i=1}^n \log(1 + e^{-y_i(w_1x_{i1} + w_2x_{i2} + \dots + w_mx_{im} + b)}) + \lambda \|w\|^2$$

Gaussian prior:

$$\operatorname{argmin}_{w,b} \sum_{i=1}^n \log(1 + e^{-y_i(w_1x_{i1} + w_2x_{i2} + \dots + w_mx_{im} + b)}) + \frac{1}{2\sigma^2} \|w\|^2$$

Does the  $\lambda$  make sense?       $\lambda = \frac{1}{2\sigma^2}$

### Regularization/prior

---


L2 regularization:

$$\operatorname{argmin}_{w,b} \sum_{i=1}^n \log(1 + e^{-y_i(w_1x_{i1} + w_2x_{i2} + \dots + w_mx_{im} + b)}) + \lambda \|w\|^2$$

Gaussian prior:

$$\operatorname{argmin}_{w,b} \sum_{i=1}^n \log(1 + e^{-y_i(w_1x_{i1} + w_2x_{i2} + \dots + w_mx_{im} + b)}) + \frac{1}{2\sigma^2} \|w\|^2$$

$$\lambda = \frac{1}{2\sigma^2}$$




## Regularization/prior

L1 regularization:

$$\operatorname{argmin}_{w,b} \sum_{i=1}^n \log(1 + e^{-y_i(w_1x_{i1} + w_2x_{i2} + \dots + w_mx_{im} + b)}) + \lambda \|w\|$$

Laplacian prior:



## Regularization/prior

L1 regularization:

$$\operatorname{argmin}_{w,b} \sum_{i=1}^n \log(1 + e^{-y_i(w_1x_{i1} + w_2x_{i2} + \dots + w_mx_{im} + b)}) + \lambda \|w\|$$

Laplacian prior:


$$\operatorname{argmin}_{w,b} \sum_{i=1}^n \log(1 + e^{-y_i(w_1x_{i1} + w_2x_{i2} + \dots + w_mx_{im} + b)}) + \frac{1}{\sigma} \|w\|$$

$$\lambda = \frac{1}{\sigma}$$

## L1 vs. L2

L1 = Laplacian prior

L2 = Gaussian prior




## Logistic regression


Why is it called logistic regression?  
It is a classifier??

$$\log \frac{P(1|x_1, x_2, \dots, x_m)}{1 - P(1|x_1, x_2, \dots, x_m)} = w_1x_1 + w_2x_2 + \dots + w_mx_m + b$$

## A digression: regression vs. classification



## linear regression




Given some points, find the **line** that best fits/explains the data

Our model is a line, i.e. we're assuming a linear relationship between the feature and the label value

$$h(y) = w_1 x_1 + b$$

How can we find this line?

## Linear regression



Learn a line  $h$  that minimizes some loss/error function:

$$error(h) = ?$$

Sum of the individual errors:

$$error(h) = \sum_{i=1}^n |y_i - h(f_i)|$$

0/1 loss!

## Error minimization

How do we find the minimum of an equation?


$$error(h) = \sum_{i=1}^n |y_i - h(f_i)|$$

Take the derivative, set to 0 and solve (going to be a min or a max)


Any problems here?

Ideas?

## Linear regression



## Linear regression



## Linear regression

We'd like to *minimize* the error

Find  $w_1$  and  $w_0$  such that the error is minimized

$$\text{error}(h) = \sum_{i=1}^n (y_i - (w_1 f_i + w_0))^2$$

We can solve this in closed form

## Multiple linear regression

If we have  $m$  features, then we have a line in  $m$  dimensions

$$h(\vec{f}) = w_0 + w_1 f_1 + w_2 f_2 + \dots + w_m f_m$$

weights

## Multiple linear regression

We can still calculate the squared error like before


$$h(\vec{f}) = w_0 + w_1 f_1 + w_2 f_2 + \dots + w_m f_m$$

$$error(h) = \sum_{i=1}^n (y_i - (w_0 + w_1 f_{i1} + w_2 f_{i2} + \dots + w_m f_{im}))^2$$

Still can solve this exactly!


## Logistic function

$$\text{logistic} = \frac{1}{1 + e^{-x}}$$



## Logistic regression

Find the best fit of the data based on a logistic



## Basic steps for probabilistic modeling

### Probabilistic models

Step 1: pick a model

Which model do we use, i.e. how do we calculate  $p(\text{feature}, \text{label})$ ?

Step 2: figure out how to estimate the probabilities for the model

How do train the model, i.e. how to we we **estimate the probabilities** for the model?

Step 3 (optional): deal with overfitting

How do we deal with overfitting?

## Probabilistic models summarized

Two classification models:

- ▣ Naïve Bayes (models **joint** distribution)
- ▣ Logistic Regression (models **conditional** distribution)
  - In practice this tends to work better if all you want to do is classify

Priors/smoothing/regularization

- ▣ Important for both models
- ▣ In theory: allow us to impart some prior knowledge
- ▣ In practice: avoids overfitting and often tune on development data